



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Mosquito-proof ward for isolation of yellow-fever patients.

Assistant Surgeon Pierce reports, February 27, through Surgeon Perry, as follows:

I have the honor to report that the authorities that control the San Tomas Hospital in this city have made one of the wards mosquito-proof, and intend to isolate in this ward all cases of yellow fever that may be sent to the hospital in the future.

The ward so prepared is 40 feet long by 20 broad; it has two double doors and no windows. Around the top of the wall is an open space 2 feet broad, which gives the only ventilation. The doors have mosquito-screen vestibules.

Since the completion of the ward no cases have been treated in the hospital, nor have any cases been reported in the city.

Foreign and insular statistical reports of countries and cities—Yearly and monthly.

AFRICA—*Sierra Leone*.—Two weeks ended February 12, 1904. Estimated population, 40,000. Total number of deaths 48. No contagious diseases.

BRAZIL—*Pernambuco*.—Two weeks ended January 31, 1904. Estimated population 200,000. Total number of deaths 363, including enteric fever 1, whooping cough 1, smallpox 28, and 42 from phthisis pulmonalis.

CANADA—*Ontario—Hamilton*.—Month of February, 1904. Estimated population, 57,000. Total number of deaths 102, including diphtheria 3 and 6 from tuberculosis.

GERMANY—*Hanover*.—Month of December, 1903. Estimated population, 250,408. Total number of deaths 309, including 14 from infectious diseases.

Strasburg.—Month of December, 1903. Estimated population, 159,006. Total number of deaths 250, including diphtheria 4, enteric fever 4, measles 4, whooping cough 2, and 43 from tuberculosis.

GIBALTAR.—Two weeks ended February 28, 1904. Estimated population, 27,460. Total number of deaths, 17. No deaths from contagious diseases.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended February 20, 1904, correspond to an annual rate of 18.5 per 1,000 of the aggregate population, which is estimated at 15,271,425.

London.—One thousand five hundred and eighty-nine deaths were registered during the week, including measles 42, scarlet fever 9, diphtheria 17, whooping cough 55, enteric fever 2, smallpox 1, and diarrhea 16. The deaths from all causes correspond to an annual rate of 17.8 per 1,000. In Greater London 2,135 deaths were registered. In the "outer ring" the deaths included 3 from diphtheria, 11 from measles, and 9 from whooping cough.